STEWART
TALENT

HEIGHT:5'9" BUST:34" CUP:C WAIST:25.5" HIPS:36" DRESS:2 SHOE SIZE:6.5 HAIR:DARK BROWN EYES:BROWN

HEIGHT:5'9" BUST:34" CUP:C WAIST:25.5" HIPS:36" DRESS:2 SHOE SIZE:6.5 HAIR:DARK BROWN EYES:BROWN



STEWART
TALENT

HEIGHT:5'9" BUST:34" CUP:C WAIST:25.5" HIPS:36" DRESS:2 SHOE SIZE:6.5 HAIR:DARK BROWN EYES:BROWN

STEWART
TALENT

HEIGHT:5'9" BUST:34" CUP:C WAIST:25.5" HIPS:36" DRESS:2 SHOE SIZE:6.5 HAIR:DARK BROWN EYES:BROWN



CLASS 1 STRENGTH

with the Cooling Sensation of L-Menthol and the Emollience of Mineral Oil[1,7]

INACTIVE INGREDIENTS[7]

GLYCERYL OLEATE
(Skin conditioner)

ISOPROPYL ALCOHOL
(Solvent)

ISOPROPYL MYRISTATE
(Penetration enhancer/Co-solvent)

L-MENTHOL
(Cooling sensation agent/ Enhancer)

MINERAL OIL
(Oleaginous vehicle/Emollient)

PROPYLENE GLYCOL-FREE[2] | SODIUM LAURYL SULFATE FREE[2] | PARABEN-FREE[2]

Topicort® Topical Spray is a topical corticosteroid indicated for the treatment of plaque psoriasis in patients 18 years of age or older.

- Local adverse reactions may be more likely to occur with occlusive use, prolonged use or use of higher potency corticosteroids. Reactions may include atrophy, striae, telangiectasias, burning, itching, irritation, dryness, folliculitis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, secondary infection, and miliaria. Some local reactions may be irreversible.
- Safety and effectiveness of Topicort® Topical Spray in patients younger than 18 years of age have not been studied; therefore use in pediatric patients is not recommended.

0.25% SPRAY Topicort® (desoximetasone) Topical Spray 0.25%

CLASS 1 STRENGTH WITH A WELL-TOLERATED SPRAY FORMULATION[1,2]

TOUGH & WELL-TOLERATED

Topicort® Topical Spray is a topical corticosteroid indicated for the treatment of plaque psoriasis in patients 18 years of age or older.

Important Safety Information

- Topicort® Topical Spray is a topical corticosteroid that has been shown to suppress the hypothalamic-pituitary-adrenal (HPA) axis.
- Systemic absorption of topical corticosteroids can produce reversible HPA axis suppression with the potential for glucocorticosteroid insufficiency. This may occur during treatment or upon withdrawal of the topical corticosteroid.
- Because of the potential for systemic absorption, use of topical corticosteroids may require that patients be periodically evaluated for HPA axis suppression.
- Local adverse reactions may be more likely to occur with occlusive use, prolonged use or use of higher potency corticosteroids. Reactions may include atrophy, striae, telangiectasias, burning, itching, irritation, dryness, folliculitis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, secondary infection, and miliaria. Some local reactions may be irreversible.
- Safety and effectiveness of Topicort® Topical Spray in patients younger than 18 years of age have not been studied; therefore use in pediatric patients is not recommended.

Before prescribing, please see enclosed Full Prescribing Information.

0.25% SPRAY Topicort® (desoximetasone) Topical Spray 0.25%

STEWART
TALENT

HEIGHT:5'9" BUST:34" CUP:C WAIST:25.5" HIPS:36" DRESS:2 SHOE SIZE:6.5 HAIR:DARK BROWN EYES:BROWN